



Peripheral Arterial Disease in Diabetes Mellitus and Physiotherapy Management: A Narrative Review

Chew Wai Hoong¹, Mah Kai Xuan^{2*}

¹ Department of Physiotherapy, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, Malaysia

² Medical Rehabilitation Unit, Tung Shin Hospital, Malaysia

ARTICLE INFO

ABSTRACT

Article history:

Received 13 October 2024

Received in revised form 1 November 2024

Accepted 20 November 2024

Available online 15 December 2024

Keywords:

Peripheral arterial disease; diabetes mellitus; physiotherapy management

Diabetes Mellitus is prevalent and contributed to high mortality rate. The hyperglycemia in diabetic individual leads to low-grade inflammation, vasoconstriction, arterial stiffness, endothelial dysfunction and thrombotic abnormalities. These placing them at higher risk of developing peripheral arterial disease which compromised the blood perfusion, particularly the lower extremity. Despite the development of peripheral arterial disease in Diabetes Mellitus can leads to wide array of complications and leaves the individual with great disability, however, various physiotherapy interventions have been supported by previous literature to be able to improve the functional outcomes and quality of life. This first part of this review paper focused on the pathophysiology of peripheral arterial disease in Diabetes Mellitus, while the second part focused on the various physiotherapy interventions for improving the functions and outcomes in diabetic individual with peripheral arterial disease.

1. Introduction

According to the International Diabetes Federation, diabetes is affecting approximately 635 million people worldwide and has contributed to more than 6 million deaths in 2021. Although the disease incidence has begun to decline in some nations, diabetes prevalence has climbed over the past few decades in the majority of other developed and developing nations especially the Southeast Asian Countries. South-East Asia is among the regions with the highest number of people with diabetes, contributing to about 90 million and the figure is projected to escalate to 151 million by 2045. The advances in the management of diabetes mellitus have successfully increased the life expectancies, however, diabetes continues to pose a considerable burden on millions of people living with diabetes mellitus. The global age-standardised disability-adjusted-life-years (DALYs) linked with diabetes in 2019 was 801.5 per 100,000, an increase of nearly 30% when compared to 1990 [32]. The healthcare cost associated with diabetes is tremendous. Globally, it has caused at least USD 966

* Corresponding author.

E-mail address: mahkaixuan97@gmail.com

<https://doi.org/10.37934/jhqol.4.1.1824>

billion dollars, more than 300% increase over the last 15 years (International Diabetes Federation, 2021) [15]. In Malaysia, it is estimated that about 20% of its adult population was having diabetes and the estimated total healthcare cost was approximately USD 600 million [10]. Spending on managing diabetes and its complications constitutes one of the direct medical expenses associated with the disease. These expenses cover long-term care, medication, and medical supplies in addition to outpatient, emergency, and inpatient hospital care.

Diabetes mellitus has been known to cause a wide array of complications including peripheral arterial disease. Previous literature suggested that Diabetes Mellitus is associated with greater severity and more diffuse peripheral arterial disease as well as lower extremity complications relative to non-diabetics, which may increase the risk for lower limb amputation four to five times higher [24]. Studies reported that chronic hyperglycemia is often being observed in diabetic individual, which leads to low-grade inflammation, vasoconstriction, arterial stiffness, endothelial dysfunction and thrombotic abnormalities [24,36]. All of which would affect and reduce the blood circulation to the limb, negatively impacting the perfusion to the lower limb. As a result, this would lead to an increased risk for amputation, cardiovascular disease, stroke and risk of death. In fact, Criqui *et al.*, [8] has reported that lower extremity arterial disease often accompanied with intermittent claudication and ulcer and is one of the main risk factors for impaired physical function and reduced physical activity.

2. Peripheral Arterial Disease in Diabetes Mellitus

Peripheral arterial disease is a form of atherosclerotic disease characterised by the occlusion of the blood vessels, compromising the blood perfusion. It is known that Diabetes Mellitus increases the risk of peripheral arterial disease and also accelerates the disease progression and severity. Previous literature has found that people with Diabetes Mellitus are at 2 to 4 times greater risk of developing peripheral arterial disease when compared to their counterpart [4]. The typical presentation of peripheral arterial disease is intermittent claudication, although some may be asymptomatic. Intermittent claudication usually manifested as cramping, numbness or burning sensation at the lower extremity which was brought upon by physical activity and relieved by rest [8]. The most severe form of peripheral arterial disease is chronic limb-threatening ischaemia. Diabetic individuals with chronic limb-threatening ischaemia often complain of diffuse burning or aching pain at the foot and leg, which somewhat improved with dependent positioning of the limb [3]. Tissue loss is a common consequence seen in diabetic individuals with peripheral arterial disease. The compromised blood perfusion at the lower extremity increases the likelihood of ulceration and infection, which eventually would lead to amputation when the tissue damage is irreversible.

In Diabetes Mellitus, the overproduction of advanced glycation end products has long been proposed as the principal mechanism for causing peripheral arterial disease [35,38]. The overproduction of advanced glycation end products (AGEs) could contribute to the development of peripheral arterial disease via 2 main pathologic pathways: 1) abnormal cross-linking of proteins and 2) cellular stress response by interaction with the receptors for advanced glycation end products [9]. Several AGEs, especially pentosidine which formed as a result of non-enzymatic glycation, played a significant role in the development of angiopathic complications in individuals with Diabetes Mellitus [16]. In the study done by Kerkeni *et al.*, [18] which investigated 200 subjects with Diabetes Mellitus, the authors reported that pentosidine level was found to be significantly higher in diabetic subjects and it is positively associated with microvascular complications in Diabetes Mellitus. Pentosidine aggravates the cross-linking formation between and within collagen and elastin proteins which adversely affect the vessel's distensibility and contribute to increased arterial stiffness [34,35].

On the other hand, the AGEs could also lead to excessive cellular stress responses via its engagement with the receptors for advanced glycation end products (RAGE). RAGE is a signal-transducing receptor for AGEs and expressed in various cell membranes including endothelial and smooth muscles [38]. Previous studies have reported there is increased expression of RAGE in people with Diabetes Mellitus [2,30]. The binding of AGEs to RAGE promotes the activation of nuclear factor-kappa B (NF- κ B) which up-regulates the release of pro-inflammatory cytokines leading to endothelial dysfunction [9]. As a result, it induces structural as well as functional vascular abnormalities, altering the reactivity and relaxation of blood vessels which stimulate atherosclerosis [24]. Along with the arterial stiffness due to the aggravated cross-linking of proteins, they contribute to the pathogenesis of peripheral arterial disease in diabetic individuals, compromising the blood perfusion to the extremity.

3. Physiotherapy Interventions for Peripheral Arterial Disease

3.1 Resistance Training

A recent randomised controlled trial by Alni and Nikookheslat [1] investigated the effectiveness of resistance training in diabetic individuals with peripheral vascular disease. The resistance training consists of various upper body, lower body and trunk exercises performed at 8-12 repetitions for two to three sets. Significant improvement was noted in the ankle-brachial index after 12 weeks of intervention and the improvement was significantly better when compared to the aerobic exercise group and control group. In Diabetes Mellitus, the overproduction of AGEs lead to dysfunction of endothelial cell as well as smooth muscle in the arteries [24,38]. In addition, the hyperglycemia also causes increased in blood viscosity and platelet aggregation in diabetic individuals which further worsen the blood flow and perfusion [12,23]. The clinical guideline by American Heart Association/American College of Cardiology has recommended the prescription of resistance exercise in patients with Diabetes Mellitus [11]. Resistance training is beneficial in increasing the muscle mass and strength which could help to improve the glucose metabolism [13]. Moreover, previous literature has also reported that exercise is beneficial in improving the concentration of endothelin-1 and nitric oxide which are essential in regulating the arterial function [25,33]. A systematic review and meta-analysis by Liu *et al.*, [20] which included 24 trials comprising more than 900 subjects has also reported that resistance training, particularly when performed at high intensity, has significant effect in decreasing the HbA1c level in diabetic individuals. This could be another plausible reason as to how resistance training could help in improving peripheral arterial disease in people with Diabetes Mellitus as HbA1c has been reported previously that it is significantly associated with vascular complications [31].

3.2 Walking Exercise

In people with Diabetes Mellitus, previous literature has reported the population often exhibit deterioration in their walking performance [19,37]. Declination in muscle strength, sarcopenia, neuropathy and intermittent claudication associated with peripheral artery disease are among the factors affecting their walking performance [5,7,17,29]. In the experimental study by McDermott *et al.*, [21], the authors have investigated the effectiveness of a 6-month supervised treadmill walking on improving walking distance. It was found that supervised treadmill walking did able to significantly improve the walking distance by 6-month follow up, however, the improvement was not maintained when being further followed up at 12-month. Contrary to supervised treadmill, Collins *et al.*, [6] investigated the effects of a home-based walking intervention in diabetic population with peripheral

arterial disease. At 6-month follow up, it was being reported that the walking speed and quality of life demonstrated significant improvement although no significant effect was observed on the subjects' walking distance. Regular walking exercise has been shown that it has positive effects on body weight, muscle strength, blood pressure, skeletal muscle oxygenation as well as glycemic control which could help to improve the walking performance in diabetic individuals with peripheral artery disease [22,27]. The exercise therapy guidelines by American Heart Association has also provided class I recommendation for the prescription of supervised exercise to improve the functional status and leg symptoms in peripheral artery disease individuals with claudication [11].

3.2 Buerger-Allen Exercise

Buerger-Allen exercise is theorised to improve the lower extremity blood circulation, aid in the removal of blood that had become stagnant as well as establishing collateral circulation to the ischemic area. This mechanism is achieved by utilising the effect of gravity and alternatively fills and empties the blood vessel, thus improves the lower limbs circulation of the individual. A study done by Radhika *et al.*, [28] in 2020 evaluated the effect on Buerger-Allen exercise in improving the lower limb perfusion among Diabetes Mellitus population and it has been reported a nearly twofold improvement in the lower limb perfusion as indicated by ankle-brachial index post-intervention. Hafid *et al.*, [14] conducted an experimental study in 2020 to see the effect of Buerger-Allen exercise on Type 2 Diabetes Mellitus population. In this particular study, a significant increase in the ABI value in the peripheral circulation was documented. These outcomes suggest the use of this strategy, Buerger-Allen exercise is effective in improving the lower extremity blood circulation in Diabetes Mellitus population as theorised. Similar outcomes were seen in the study done by Patidar *et al.*, [26] in 2018 as well. Another similar research was done in 2018 by Zahran *et al.*, [39] on 48 adult patients with type 2 Diabetes Mellitus who visited a specialised medical centre for six months. According to the authors, Buerger-Allen exercise was able to significantly improve the lower limb perfusion both immediately post-intervention and even four weeks later when the subjects were being monitored again.

4. Conclusion

Diabetes is amongst the biggest global public health issues, having a significant negative impact on both socioeconomic development and public health globally. The disease, if not managed well, often leading to high disability and substantial healthcare cost. The overproduction of AGEs seen in Diabetes Mellitus causes abnormal cross-linking of proteins and cellular stress response which adversely affecting the structural and functionality of artery, giving rise to peripheral artery disease. Peripheral artery disease characterised by occlusion of the vessel and the most common presentation is intermittent claudication. Previous studies have provided evidences in supporting physiotherapy interventions such as resistance training, walking intervention as well as Buerger-Allen exercise in improving the leg symptoms and function in diabetic individuals with peripheral artery disease.

References

- [1] Alni, Masoud Abdollahpour, and Saeid Dabbagh Nikookheslat. "The effect of 12 weeks aerobic, resistance and combined trainings on peripheral vascular disease in type 2 diabetes with peripheral neuropathy in men." *Obesity Medicine* 34 (2022): 100439. <https://doi.org/10.1016/j.obmed.2022.100439>
- [2] Barlovic, Drazenka Pongrac, Aino Soro-Paavonen, and Karin AM Jandeleit-Dahm. "RAGE biology, atherosclerosis and diabetes." *Clinical science* 121, no. 2 (2011): 43-55. <https://doi.org/10.1042/CS20100501>

- [3] Barnes, J. Aaron, Mark A. Eid, Mark A. Creager, and Philip P. Goodney. "Epidemiology and risk of amputation in patients with diabetes mellitus and peripheral artery disease." *Arteriosclerosis, thrombosis, and vascular biology* 40, no. 8 (2020): 1808-1817. <https://doi.org/10.1161/ATVBAHA.120.314595>
- [4] Beckman, Joshua A., Mark A. Creager, and Peter Libby. "Diabetes and atherosclerosis: epidemiology, pathophysiology, and management." *Jama* 287, no. 19 (2002): 2570-2581. <https://doi.org/10.1001/jama.287.19.2570>
- [5] Chung, Seung Min, Jun Sung Moon, and Min Cheol Chang. "Prevalence of sarcopenia and its association with diabetes: a meta-analysis of community-dwelling Asian population." *Frontiers in medicine* 8 (2021): 681232. <https://doi.org/10.3389/fmed.2021.681232>
- [6] Collins, Tracie C., Scott Lunos, Teri Carlson, Kimberly Henderson, Marissa Lightbourne, Brittney Nelson, and James S. Hodges. "Effects of a home-based walking intervention on mobility and quality of life in people with diabetes and peripheral arterial disease: a randomized controlled trial." *Diabetes care* 34, no. 10 (2011): 2174-2179. <https://doi.org/10.2337/dc10-2399>
- [7] Correia, Marilia Almeida, Gustavo Oliveira Silva, Paulo Longano, Ivani Credidio Trombetta, Fernanda Consolim-Colombo, Pedro Puech-Leão, Nelson Wolosker, Gabriel Grizzo Cucato, and Raphael Mendes Ritti-Dias. "In peripheral artery disease, diabetes is associated with reduced physical activity level and physical function and impaired cardiac autonomic control: A cross-sectional study." *Annals of Physical and Rehabilitation Medicine* 64, no. 2 (2021): 101365. <https://doi.org/10.1016/j.rehab.2020.01.006>
- [8] Criqui, Michael H., Kunihiro Matsushita, Victor Aboyans, Connie N. Hess, Caitlin W. Hicks, Tak W. Kwan, Mary M. McDermott, Sanjay Misra, and Francisco Ujueta. "Lower extremity peripheral artery disease: contemporary epidemiology, management gaps, and future directions: a scientific statement from the American Heart Association." *Circulation* 144, no. 9 (2021): e171-e191. <https://doi.org/10.1161/CIR.0000000000001005>
- [9] de Vos, Lianne C., Joop D. Lefrandt, Robin PF Dullaart, Clark J. Zeebregts, and Andries J. Smit. "Advanced glycation end products: An emerging biomarker for adverse outcome in patients with peripheral artery disease." *Atherosclerosis* 254 (2016): 291-299. <https://doi.org/10.1016/j.atherosclerosis.2016.10.012>
- [10] Ganasegeran, Kurubaran, Chee Peng Hor, Mohd Fadzly Amar Jamil, Hong Chuan Loh, Juliana Mohd Noor, Norshahida Abdul Hamid, Purnima Devi Suppiah, Mohd Rizal Abdul Manaf, Alan Swee Hock Ch'ng, and Irene Looi. "A systematic review of the economic burden of type 2 diabetes in Malaysia." *International journal of environmental research and public health* 17, no. 16 (2020): 5723. <https://doi.org/10.3390/ijerph17165723>
- [11] Drachman, D. E., L. A. Fleisher, F. G. R. Fowkes, N. M. Hamburg, S. Kinlay, R. Lookstein, S. Misra et al. "2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease." (2016).
- [12] Gresele, Paolo, Stefania Marzotti, Giuseppe Guglielmini, Stefania Momi, Silvia Giannini, Pietro Minuz, Paola Lucidi, and Geremia B. Bolli. "Hyperglycemia-induced platelet activation in type 2 diabetes is resistant to aspirin but not to a nitric oxide-donating agent." *Diabetes care* 33, no. 6 (2010): 1262-1268. <https://doi.org/10.2337/dc09-2013>
- [13] Hap, K., Biernat, K. and Konieczny, G., 2021. Patients with Diabetes Complicated by Peripheral Artery Disease: the Current State of Knowledge on Physiotherapy Interventions. *Journal of Diabetes Research*, 2021, pp.1-12. <https://doi.org/10.1155/2021/5122494>
- [14] Hafid, Muh Anwar, Ilhamsyah Ilhamsyah, and Mutmainnah Sari. "The effect of buerger allen exercise on the lower extremity peripheral circulation in type 2 diabetic patients." *STRADA Jurnal Ilmiah Kesehatan* 10, no. 1 (2021): 612-621.
- [15] Federation, Intenational Diabetes. "IDF diabetes atlas, tenth." *International Diabetes* (2021).
- [16] Jaishankar, Sarath Kumar Jaganathan, Amrutha Murali, And Amrin Marzooka. "Assessment of Pentosidine as a Marker for Advanced Glycation End Products in Type 2 Diabetes Mellitus Patients." *Journal of Clinical & Diagnostic Research* 14, no. 2 (2020). <https://doi.org/10.7860/JCDR/2020/42745.13488>
- [17] Kennedy, L., 2007. Decreased Muscle Strength and Quality in Older Adults With Type 2 Diabetes: The Health, Aging, and Body Composition Study. *Yearbook of Endocrinology*, 2007, pp.25-26. [https://doi.org/10.1016/S0084-3741\(08\)70018-0](https://doi.org/10.1016/S0084-3741(08)70018-0)
- [18] Kerkeni, Mohsen, Amel Saïdi, Hassan Bouzidi, Ahmed Letaief, Salim Ben Yahia, and Mohamed Hammami. "Pentosidine as a biomarker for microvascular complications in type 2 diabetic patients." *Diabetes and Vascular Disease Research* 10, no. 3 (2013): 239-245. <https://doi.org/10.1177/1479164112460253>
- [19] Kuziemski, Krzysztof, Wojciech Słomiński, and Ewa Jassem. "Impact of diabetes mellitus on functional exercise capacity and pulmonary functions in patients with diabetes and healthy persons." *BMC endocrine disorders* 19 (2019): 1-8. <https://doi.org/10.1186/s12902-018-0328-1>
- [20] Liu, Yubo, Weibing Ye, Qian Chen, Yong Zhang, Chia-Hua Kuo, and Mallikarjuna Korivi. "Resistance exercise intensity is correlated with attenuation of HbA1c and insulin in patients with type 2 diabetes: a systematic review and meta-analysis." *International journal of environmental research and public health* 16, no. 1 (2019): 140. <https://doi.org/10.3390/ijerph16010140>

- [21] McDermott, Mary M., Melina R. Kibbe, Jack M. Guralnik, Luigi Ferrucci, Michael H. Criqui, Kathryn Domanchuk, Lu Tian et al. "Durability of benefits from supervised treadmill exercise in people with peripheral artery disease." *Journal of the American Heart Association* 8, no. 1 (2019): e009380. <https://doi.org/10.1161/JAHA.118.009380>
- [22] Moghetti, P., S. Balducci, L. Guidetti, P. Mazzuca, E. Rossi, and F. Schena. "Walking for subjects with type 2 diabetes: a systematic review and joint AMD/SID/SISMES evidence-based practical guideline." *Nutrition, Metabolism and Cardiovascular Diseases* 30, no. 11 (2020): 1882-1898. <https://doi.org/10.1016/j.numecd.2020.08.021>
- [23] Mushtaq, Mazhar, Muhammad Abdul Mateen, and Uh-Hyun Kim. "Hyperglycemia associated blood viscosity can be a nexus stimuli." *Clinical hemorheology and microcirculation* 71, no. 1 (2019): 103-112. <https://doi.org/10.3233/CH-180426>
- [24] Nativel, Mathilde, Louis Potier, Laure Alexandre, Laurence Baillet-Blanco, Eric Ducasse, Gilberto Velho, Michel Marre, Ronan Roussel, Vincent Rigalleau, and Kamel Mohammedi. "Lower extremity arterial disease in patients with diabetes: a contemporary narrative review." *Cardiovascular diabetology* 17 (2018): 1-14. <https://doi.org/10.1186/s12933-018-0781-1>
- [25] Otsuki, Takeshi, Fumiko Nakamura, and Asako Zempo-Miyaki. "Nitric oxide and decreases in resistance exercise blood pressure with aerobic exercise training in older individuals." *Frontiers in Physiology* 10 (2019): 1204. <https://doi.org/10.3389/fphys.2019.01204>
- [26] Patidar, V. "A study to assess the effectiveness of burger allen exercise on improving peripheral circulation among type 2 Diabetes Mellitus patients in selected hospitals of Nadiad city." *Clinical Practice* 15, no. 6 (2018). <https://doi.org/10.4172/clinical-practice.1000425>
- [27] Qiu, Shanhu, Xue Cai, Uwe Schumann, Martina Velders, Zilin Sun, and Jürgen Michael Steinacker. "Impact of walking on glycemic control and other cardiovascular risk factors in type 2 diabetes: a meta-analysis." *PloS one* 9, no. 10 (2014): e109767. <https://doi.org/10.1371/journal.pone.0109767>
- [28] Radhika, Jinna, Geetha Poomalai, SiralaJagadeesh Nalini, and Ramanathan Revathi. "Effectiveness of buerger-allen exercise on lower extremity perfusion and peripheral neuropathy symptoms among patients with diabetes mellitus." *Iranian journal of nursing and midwifery research* 25, no. 4 (2020): 291-295. https://doi.org/10.4103/ijnmr.IJNMR_63_19
- [29] Rahimi, Mohammad, Payam Saadat, Seyed reza Hosseini, Mohammad Ali Bayani, and Ali Bijani. "Muscle strength in diabetics compared to non-diabetic elderly subjects: A cross sectional and case-control study." *Caspian Journal of Internal Medicine* 10, no. 3 (2019): 265.
- [30] Ramasamy, Ravichandran, Shi Fang Yan, and Ann Marie Schmidt. "Receptor for AGE (RAGE): signaling mechanisms in the pathogenesis of diabetes and its complications." *Annals of the New York Academy of Sciences* 1243, no. 1 (2011): 88-102. <https://doi.org/10.1111/j.1749-6632.2011.06320.x>
- [31] Rosenson, R. S., Paola Fioretto, and P. M. Dodson. "Does microvascular disease predict macrovascular events in type 2 diabetes?." *Atherosclerosis* 218, no. 1 (2011): 13-18. <https://doi.org/10.1016/j.atherosclerosis.2011.06.029>
- [32] Safiri, Saeid, Nahid Karamzad, Jay S. Kaufman, Arielle Wilder Bell, Seyed Aria Nejadghaderi, Mark JM Sullman, Maziar Moradi-Lakeh, Gary Collins, and Ali-Asghar Kolahi. "Prevalence, deaths and disability-adjusted-life-years (DALYs) due to type 2 diabetes and its attributable risk factors in 204 countries and territories, 1990-2019: results from the global burden of disease study 2019." *Frontiers in endocrinology* 13 (2022): 838027. <https://doi.org/10.3389/fendo.2022.838027>
- [33] Sanian, Hassan, Hassan Matinhomae, and Maghsoud Peeri. "The effect of circuit resistance training on plasma concentration of endothelin-1, nitric oxide and vascular diameter in elderly men." *Journal of Clinical Research in Paramedical Sciences* 8, no. 1 (2019). <https://doi.org/10.5812/jcrps.80932>
- [34] Sell, David R., and Vincent M. Monnier. "Molecular basis of arterial stiffening: role of glycation—a mini-review." *Gerontology* 58, no. 3 (2012): 227-237. <https://doi.org/10.1159/000334668>
- [35] Soyoye, David Olubukunmi, Olugbenga Olusola Abiodun, Rosemary Temidayo Ikem, Babatope Ayodeji Kolawole, and Anthony Olubunmi Akintomide. "Diabetes and peripheral artery disease: A review." *World journal of diabetes* 12, no. 6 (2021): 827. <https://doi.org/10.4239/wjd.v12.i6.827>
- [36] Williams, Dean T., Keith G. Harding, and Patricia Price. "An evaluation of the efficacy of methods used in screening for lower-limb arterial disease in diabetes." *Diabetes care* 28, no. 9 (2005): 2206-2210. <https://doi.org/10.2337/diacare.28.9.2206>
- [37] Wu, Ran, Yan Zhang, Jiao-Jiao Bai, Jiao Sun, Zhi-Jun Bao, and Zheng Wang. "Impact of lower limb muscle strength on walking function beyond aging and diabetes." *Journal of International Medical Research* 48, no. 6 (2020): 0300060520928826. <https://doi.org/10.1177/0300060520928826>

- [38] Yang, Shao-ling, Lv-yun Zhu, Rui Han, Lei-lei Sun, Jun-xia Li, and Jing-tao Dou. "Pathophysiology of peripheral arterial disease in diabetes mellitus: 糖尿病周围动脉疾病的病理生理机制." *Journal of diabetes* 9, no. 2 (2017): 133-140. <https://doi.org/10.1111/1753-0407.12474>
- [39] Zahran, Walaa, Amira Hassanen, Madiha Nabih, and Fady Kyrillos. "Effect of Buerger Allen exercise on lower limb perfusion among patients with type 2 diabetes mellitus." *Mansoura Nursing Journal* 5, no. 1 (2018): 101-111. <https://doi.org/10.21608/mnj.2018.150616>